

## iBalance® Primary TKA Outcomes

### PURPOSE

To report the clinical outcomes for pain, function and quality of life for patients who have undergone primary TKA with the iBalance prosthesis.

### METHODS


The inclusion criteria for this analysis were patients enrolled in the Surgical Outcomes System™ global registry who underwent primary TKA with the iBalance prosthesis. Standard patient-reported outcomes questionnaires for VAS, KOOS Jr and Oxford Knee Score were administered at standard time points postoperatively. Results were reported from presurgery out to 4 years postsurgery. The number of patients included per time point is shown to the right.

Time Point	# of Compliant Patients/ Total # of Patients
Presurgery	1142/1166
1 year	815/1013
2 years	689/923
3 years	424/691
4 years	201/363


### TREND CONCLUSION

Based on these results, the pain, function and quality-of-life scores for patients undergoing primary TKA with the iBalance prosthesis trend towards favorable outcomes. However, no claims can be made on the potential of these results without further statistical analysis to determine if there is any statistical significance.


### RESULTS



VAS



KOOS Jr



Oxford Knee Score



<b>Time Point</b>	<b>iBalance® Prosthesis Avg +/- STD VAS</b>
Presurgery	5.9 +/- 2.5
1 year	1.5 +/- 2.0
2 years	1.3 +/- 2.0
3 years	1.1 +/- 1.8
4 years	1.1 +/- 1.9

<b>Time Point</b>	<b>iBalance Prosthesis Avg +/- STD KOOS Jr</b>
Presurgery	44.3 +/- 13.5
1 year	77.4 +/- 16.3
2 years	81.1 +/- 16.9
3 years	82.8 +/- 17.2
4 years	83.1 +/- 17.4

<b>Time Point</b>	<b>iBalance Prosthesis Avg +/- STD Oxford Knee</b>
Presurgery	23.2 +/- 7.4
1 year	39.4 +/- 7.5
2 years	40.5 +/- 7.7
3 years	41.3 +/- 7.3
4 years	41.4 +/- 7.6